

Ethno-botanical Verification and Phytochemical Profile of Ethanolic leaves Extract of Two Medicinal Plants (*Phragmenthera capitata* and *Lantana camara*) used in Nigeria using GC-MS Technique

George, U. U.^{1*}, Mbong, E. O.², Bolarinwa K. A.³, Abiaobo, N. O.⁴

¹Department of Fisheries and Aquaculture, Faculty of Agriculture, Akwa Ibom State University, Ikot Akpaden, Akwa Ibom State, Nigeria

²Department of Biological Sciences, Ritman University, Ikot Ekpene, Akwa Ibom State, Nigeria

³Department of Education (Biology-Unit), Distance Learning Institute, University of Lagos, Nigeria.

⁴Department of Zoology, Akwa Ibom State University, Ikot Akpaden, Akwa Ibom State, Nigeria

Citation: George, U. U., Mbong, E. O., Bolarinwa, K. A., Abiaobo, N. O. (2023). Ethno-botanical Verification and Phytochemical Profile of Ethanolic leaves Extract of Two Medicinal Plants (*Phragmenthera capitata* and *Lantana camara*) used in Nigeria using GC-MS Technique. Acta Biology Forum. DOI: <https://doi.org/10.51470/ABF.2023.2.3.1>

Corresponding Author: Dr. U. U. George | E-Mail: (ubonggeorge@aksu.edu.ng)

Received 18 March 2023 | Revised 26 August 2023 | Accepted 29 September 2023 | Available Online October 15 2023

ABSTRACT

Background: The present experiment was designed to determine the bioactive constituents from extracts of leaves of *Phragmenthera capitata* and *Lantana camara*. The medicinal value of a plant species is dependent upon its various phytochemical constituents.

Methodology: The chemical compositions of the ethanolic extract of the leaves of *P. capitata* and *L. camara* were investigated using the Gas Chromatography-Mass spectrometry model GCMS-QP2010 PLUS SHIMADZU. The column used was a Perkin Elmer Elite - 5 capillary columns measuring 30m × 0.25mm with a film thickness of 0.25mm composed of 95% Dimethylpolysiloxane. The carrier gas used was Helium at a flow rate of 0.5ml/min.

Results: Fifteen (15) bioactive phytochemical compounds were identified for *P. capitata* and twenty (20) bioactive phytochemical compounds were identified for *L. camara*. The prevailing compounds in *P. capitata* were 3-H-pyrazol-3-1, Decanoic acid, n-Hexadecanoic acid, Pentanoic acid, Methyloctanoic acid, 4-octanoic acid, 15-tetraosenoic acid, oleic acid, Octadecanoic acid, Pentafluoropropionic acid, (E)-13-docosenoic acid, Vitamin E, Lupeol and Chilosyphone. Also, in *L. camara* the following compounds were identified, 3-n-hexythiolane, Docosanoic acid, 1-nonanol, 1-hexacosanol, Divalonic acid, Decanoic acid, Octanoic acid, Eicosanoic acid, Hexadecanoic acid, Nonadecanoic acid, 9-12 octadecanoic acid, 11-octadecadienoic acid, Pentadecanoic acid, Oleic acid, Methyl-Beta-D-arabinopyranoside, Acetic acid, Pentanoic acid, Heneicosanoic acid, methyl ester and 2H-pyran, 2-(7-heptadecynloxytetrahydro.

Conclusion: Based on the results of findings, it could be concluded that extract from *P. capitata* and *L. camara* may have antioxidant, antimicrobial, anti-cancer, anti-diabetic, anti-fungal and anti-convulsant activities due to the presence of secondary metabolites in the ethanolic extract.

Keywords: Ethno-botanical verification, phytochemical profile, *Phragmenthera capitata*, *Lantana camara*, GC-MS Technique, Ethanolic leaves.

1.0 INTRODUCTION

The knowledge of plants chemical constituents is key to the discovery of novel therapeutic values, hence the need for studies on plant phytochemical constituents cannot be over emphasized. All over Nigeria and the world beyond, medicinal plant use in folk medicine keeps increasing by the day even as Infectious diseases contribute significantly to high mortality rates judging from a global scale [4].

The therapeutic properties of plants have been used by mankind for centuries with zero or negligible signs of toxicity. These plants have been essential parts of indigenous traditional herbal medicine practice and their role in primary health care systems can never be underrated in developing countries including Nigeria becoming increasingly popular in the developed [35]. Research has revealed that the therapeutic values of plants

more often than not are linked to the nature and number of hosts of active principles known as phytochemicals (which include alkaloids, tannins, flavonoids, steroids, terpenoids, saponins, and glycosides) which they contain.

Lantana Camara is one such invasive alien species and is considered by IUCN as one of the world's 100 most invasive species; and among the world's 10 worst weeds. Invasion by non-native exotic plant species poses a serious threat to native plant communities and ecosystem properties, such as population dynamics and community structure [29], and alters native vegetation and causes a threat to biodiversity [29].

Mistletoe plants are hemiparasitic plants that grow on trees such as kola nut (*Cola nitida*), Avocado (*Persea americana*), Grape (*Vitis vinifera*), Orange (*Citrus sinensis*) Pear (*Pyrus communis*), etc. They are known scientifically for rich

Copyright: This is an open-access article distributed under the terms of the Creative Commons Attribution License (CC BY 4.0), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Table 1: Bioactive compounds detected from ethanolic extract of leaves of *Phragmenthera capitata*.

Peak	Retention time	% Peak Area	% Peak Height	Compound Analyzed	Molecular Weight	Molecular formula
1.	7.628	1.28	1.41	3H-Pyrazol-3-one	126	C ₆ H ₁₀ N ₂ O
2.	10.658	1.28	6.08	Decanoic acid	186	C ₁₁ H ₂₂ O ₂
3.	11.526	8.29	9.29	n-Hexadecanoic acid	256	C ₁₆ H ₃₂ O ₂
4.	12.956	0.55	2.36	Pentanoic acid	130	C ₇ H ₁₄ O ₂
5.	13.761	5.75	7.14	Methyloctanoic acid	158	C ₉ H ₁₈ O ₂
6.	15.667	0.73	2.54	4-Octenoic acid	156	C ₉ H ₁₆ O ₂
7.	17.139	4.80	16.86	Tetradecanoic acid	228	C ₁₄ H ₂₈ O ₂
8.	18.941	1.21	4.02	15-Tetracosenoic acid	380	C ₂₅ H ₄₈ O ₂
9.	20.178	29.52	23.28	Oleic acid	282	C ₁₈ H ₃₄ O ₂
10.	20.389	19.31	15.51	Octadecanoic acid	284	C ₁₈ H ₃₆ O ₂
11.	21.658	1.40	3.36	Pentafluoropropionic acid	332	C ₁₅ H ₂₅ F ₅ O ₂
12.	23.524	1.51	3.11	(E)-13-Docosenoic acid	338	C ₂₂ H ₄₂ O ₂
13.	25.156	1.17	1.96	Vitamin E	430	C ₂₉ H ₅₀ O ₂
14.	26.499	2.38	2.05	Lupeol	426	C ₃₀ H ₅₀ O
15.	27.913	0.81	1.04	Chiloscyphone	218	C ₁₅ H ₂₂ O
Total		100.0	100.0			

Table 2: Biological Activities of Phytocomponents Identified in the Ethanolic extract of *Phragmenthera capitata*

S/N	Name of the compound	Nature of Compound	Biological activity	References
1.	3H-Pyrazol-3-one	Heterocyclic organic compound	Anti-microbial, anti-fungal, anti-inflammatory, anti-cancer, anti-viral, anti-convulsant	[13]
2.	Decanoic acid	Straight chain saturated fatty acid	Antibacterial, anti-inflammatory, human metabolites	[13; 5]
3.	n-Hexadecanoic acid	Saturated fatty acid	Anti-oxidants, nematocide and pesticides	[12;33]
4.	Pentanoic acid	Organic acid	anticonvulsant	[13, 6]
5.	Methyloctanoic acid	Carboxylic acid	Anti-bacterial	[13, 5]
6.	4-Octenoic acid	Unsaturated fatty acid	Anti-bacterial, human metabolite	[13, 5]
7.	Tetradecanoic acid	Straight chain saturated fatty acid	Anti-fungal, anti-bacterial	[13, 5]
8.	15-Tetracosenoic acid	Monounsaturated Omega-9-fatty acid	Anti-bacterial	[13, 5]
9.	Oleic acid	Unsaturated fatty acid	Antibacterial	[38]
10.	Octadecanoic acid	Saturated fatty acid	Anti-inflammatory	[6]
11.	Pentafluoropropionic acid	Carboxylic acid	Efficient catalyst	[6]
12.	(E)-13-Docosenoic acid	Monounsaturated Omega-9-fatty acid	Anti-bacterial	[13]
13.	Vitamin E	Fat-soluble vitamin	Anti-oxidant	[12, 32]
14.	Lupeol	Organic compound	anti-inflammatory, anti-carcinogenic, Antibacterial, anti-oxidant, anti-diabetic	[12; 13; 23]
15.	Chiloscyphone	Organic compound	Antibacterial	[13]

3.2 GC-MS analysis of bioactive compounds identified from the fractions of ethanolic extract of *P. capitata*

The results pertaining to GC-MS analysis lead to the identification of a number of chemical constituents from the GC fractions of ethanolic extract of *L. montevidensis*. Twenty (20) bioactive compounds were identified and their, retention time, % peak area, % peak height, molecular formula, molecular weight, and percentage composition in the ethanolic extract of *L. montevidensis* is presented in (Fig. 2, Table 3). The compounds were Oleic acid (26.04 %) as the major component followed by Nonadecanoic acid (21.07 %), Methyl-.beta.-D-arabinopyranoside (12.82 %), 11-octadecanoic acid (8.04 %), 9,12-Octadecadienoic acid (5.17 %), n-Hexadecanoic acid (5.00 %), Hexadecanoic acid (4.93 %), Eicosanoic acid (4.85), Pentadecanoic acid (2.04 %), 3-n-Hexylthiolane, S,S-dioxide (1.96 %), 1-Hexacosanol acid (1.21 %), 2H-Pyran, 2-(7-heptadecyloxy)tetrahydro- (1.19 %), Octanoic acid (1.10 %), Decanoic acid (1.04 %), Decanoic acid (0.95 %), Acetic acid (0.72 %), Heneicosanoic acid, methyl ester (0.57 %), 1-Nonanol (0.55 %), Divalonic acid (0.45 %) and Pentanoic acid having (0.28 %) respectively. Major phyto-compounds obtained and their biological activities have been tabulated in (Table 4).




FIG. 2: GC-MS CHROMATOGRAM OF ETHANOLIC EXTRACT OF *L. montevidensis* LEAF

Table 3: Bioactive compounds detected from ethanolic extract of leaves of *Lantana montevidensis*

Peak	Retention time	% Peak Area	% Peak Height	Compound Analyzed	Molecular Weight	Molecular formular
1.	3.835	0.55	1.17	1-Nonanol	144	C ₉ H ₂₀ O
2.	7.713	0.45	0.56	Divalonic acid	130	C ₆ H ₁₀ O ₃
3.	10.656	1.04	3.29	Decanoic acid, methyl ester	186	C ₁₁ H ₂₂ O ₂
4.	11.524	5.00	3.97	n-Hexadecanoic acid	256	C ₁₆ H ₃₂ O ₂
5.	12.957	1.10	3.25	Octanoic acid	158	C ₉ H ₁₈ O ₂
6.	13.760	4.85	3.82	Eicosanoic acid	312	C ₂₀ H ₄₀ O ₂
7.	15.677	4.93	10.65	Hexadecanoic acid	270	C ₁₇ H ₃₄ O ₂
8.	17.111	21.07	9.81	Nonadecanoic acid	298	C ₁₉ H ₃₈ O ₂
9.	18.893	5.17	10.95	9,12-Octadecadienoic acid	294	C ₁₉ H ₃₄ O ₂
10.	18.932	8.04	13.65	11-Octadecenoic acid	296	C ₁₉ H ₃₆ O ₂
11.	19.314	2.04	5.03	Pentadecanoic acid	270	C ₁₇ H ₃₄ O ₂
12.	20.147	26.04	13.64	Oleic Acid	282	C ₁₈ H ₃₄ O ₂
13.	20.365	12.82	8.94	Methyl-.beta.-D-arabinopyranoside	164	C ₆ H ₁₂ O ₅
14.	21.650	0.72	1.65	Acetic acid	200	C ₁₂ H ₂₄ O ₂
15.	21.902	0.28	0.87	Pentanoic acid, 4-methyl-, methyl	130	C ₇ H ₁₄ O ₂
16.	23.521	1.96	2.48	3-n-Hexylthiolane, S,S-dioxide	204	C ₁₀ H ₂₀ O ₂ S
17.	23.900	0.95	3.18	Docosanoic acid, methyl ester	354	C ₂₃ H ₄₆ O ₂
18.	25.001	1.21	0.90	1-Hexacosanol	382	C ₂₆ H ₅₄ O
19.	25.611	0.57	1.59	Heneicosanoic acid, methyl ester	340	C ₂₂ H ₄₄ O ₂
20.	26.384	1.19	0.63	2H-Pyran, 2-(7-heptadecyloxy)tetrahydro-	336	C ₂₂ H ₄₀ O ₂
Total		100.0	100.0			

Table 4: Biological Activities of Phytocomponents Identified in the Ethanolic extract of *Lantana camara*

S/N	Name of the compound	Nature of Compound	Biological activity	References
1.	1-Nonanol	Straight chain fatty acid	Anti-bacterial, anti-fungal	[26; 27]
2.	Divalonic acid	Cyclic organic compound	Anti-bacterial	[26; 27]
3.	Decanoic acid	Straight chain saturated fatty acid	Antibacterial, anti-inflammatory, metabolites	[26; 27]
4.	n-Hexadecanoic acid	Saturated fatty acid	Anti-oxidants, nematocides and pesticides	[26]
5.	Octanoic acid	Saturated fatty acid	Anti-bacterial and human metabolite	[6; 33]

6.	Eicosanoic acid	Straight chain saturated fatty acid	Anti-bacterial	[6]
7.	Hexadecanoic acid	Saturated fatty acid	Anti-oxidant, nematocide and pesticides	[6; 29]
8.	Nonadecanoic acid	Straight chain saturated fatty acid	Anti-fungal	[36]
9.	9,12-Octadecadienoic acid	Poly-unsaturated omega-6- fatty acid	Anti-bacterial	[6]
10.	11-Octadecadienoic acid	Unsaturated monocarboxylic acid	Anti-bacterial	[6]
11.	Pentadecanoic acid	Straight chain saturated fatty acid	Anti-bacterial, antifungal	[27]
12.	Oleic Acid	Unsaturated fatty acid	Antibacterial	[6]
13.	Methyl-.beta.-D-arabinopyranoside	Organic compound	Antibacterial	[6]
14.	Acetic acid	Organic compound	Anti-bacterial	[6]
15.	Pentanoic acid	Organic acid	anticonvulsant	[6]
16.	3-n-Hexylthiolane	Organic compound	Antibacterial	[6]
17.	Docosanoic acid	Long-chain saturated fatty acid	moisturizers	[33]
18.	1-Hexacosanol	Saturated fatty acid	Anti-microbial, anti-bacteria,	[27; 18]
19.	Heneicosanoic acid, methyl ester	Fatty acid	Plant metabolite	[33]
20.	2H-Pyran, 2-(7-heptadecynyloxy) tetrahydro-	Organic compound	Antibacterial	[6]

4.0 DISCUSSION

The study of organic chemicals found in plants, as well as their actions, has grown in popularity. The combination of GC (best separation technique) with MS (best identification technique) made GC-MS one of the ideal techniques for quantitative analysis of volatile and semi-volatile compounds, [21].

The identified compounds in both extracts of *P. capitata* and *L. camara* showed a wide range of potent bioactivity. These phytochemicals are responsible for various pharmacological actions like antioxidants and antimicrobial activities, [37]. Among the fifteen compounds identified in *P. capitata*, 10 showed Anti-microbial activity, 4 showed Anti-inflammatory, 2 showed anti-fungal, 1 showed anti-cancer, 1 exhibited anti-viral, 2 showed anti-convulsant, 3 showed anti-oxidant and also showed activities such as anti-carcinogenic, anti-diabetic, nematocides, pesticides, metabolites and as a catalyst. In *L. camara* 20 bioactive compounds were identified, with similar biological activity as observed in *P. capitata*.

The results of the GC-MS analysis of ethanolic extract of *P. capitata* and *L. camara* are in deviant from reports of previous authors in related studies who reported lower bioactive compounds as compared to the results of the present study. [30] reported on GC-MS profiling and bioactivity prediction of compounds from *Momordica charantia* (L.) extract and identified only five (5) bioactive compounds, [20] in his studies on GC-MS analysis of *Majideazanquebarica* extract using gas chromatography and mass spectrophotometry technique identified eight (8) bioactive compounds, [10] when working on qualitative phytochemical and GC-MS analysis of some commonly consumed vegetables, identified nine (9) bioactive compounds each in *T. occidentalis* and *O. gratissimum* respectively, and reported eight (8) bioactive compounds for *T. triangulare*, [2] during their studies on the GC-MS analysis of amethanolic extract of *vernonia cinerea* identified nine (9) bioactive compounds and [34] identified nine (9) bioactive compounds when working on an extract of *Guiera senegalensis* using GC-MS analysis. Similarly, [28] reported eight (8) bioactive compounds during their studies on GC-MS analysis of methanolic extract of *H. Africana*.

The results of the present study are in agreement with the

reports of [25] who identified twenty-two (22) bioactive compounds when investigating the extract of leaves and fruits of *Trichosantheschoica* using GC-MS analysis, [7] identified 19 bioactive compounds when reporting on the GC-MS analysis of a methanolic extract of tubers of *P. tuberosa*, [8] reported 15 phyto-compounds when working on *Latana camara* and [9] identified 11 bioactive compounds when working on *Withaniasomnifera* (L.) extract using GC-MS analysis.

Phytochemicals are biologically active chemical compounds in plants that wield both medicinal and nutritional potentials [19]. This is evident by the fact that n-hexadecanoic acid, one of the major compounds in the extract of both plant species has been described as a potent antioxidant as well as a dependable anticancer agent [3]. Also, the presence of lupeol in the extract of *P. capitata* has earlier been described as having both anti-cancer and anti-inflammatory properties [23]. Oleic acid is found to have anti-bacterial activity; Vitamin E has been confirmed to have anti-oxidant properties [32].

The therapeutic potentials of bioactive compounds isolated in the extract of *P. capitata* and *L. camara* are presented in Table 2 and Table 4 with relevant citations to confirm its therapeutic activities. The results of the present study showed that extract from *P. capitata* and *L. camara* has pharmacological potentials based on the bioactive compounds isolated from the two extracts. The identified compounds possess some important biological potentials for future nematocide, pesticides, antibacterial, anti-inflammatory, anti-fungal, anti-convulsant antioxidants and metabolites. The results of the bioactive compounds obtained in *L. camara* has been confirmed by [26] when reporting on Comparative LC-LTQ-MS-MS analysis of the leaf extracts of *Lantana camara* and *Lantana montevidensis* growing in Egypt with insights into their antioxidant, anti-Inflammatory, and cytotoxic activities.

4.1 CONCLUSION

In conclusion, the major components of the ethanolic extract of *L. camara* and *Phragmenthera capitata*, as indicated in this study contains bioactive compounds of numerous biological or therapeutic importance. The fifteen and twenty phytochemical constituents identified by the GC-MS analysis of both plant

extract are therefore, medicinally valuable and possess various pharmaceutical applications. The biological activities of each of the identified phyto-components range from antimicrobial, anti-inflammatory, antioxidant, anti-cancer, anti-convulsant, and anti-fungal activities. The research findings have shown that the extract from these plants species is extensively rich in secondary metabolites. The plant's extract has a high potential for a vast number of bioactive compounds which justifies its use for various ailments by traditional practitioners. These findings have provided a scientific basis to the ethno-medical usage of the plant. However, isolation of the individual phytochemical constituents, and subjecting it to biological activity and toxicity profile will give fruitful results. Therefore, further studies isolating individual phyto-compounds to confirm its potential is recommended.

List of Abbreviations

P. capitata- *Phragmanthera capitata*

L. camara – *Latana camara*

GC – Gas Chromatography

MS – Mass Spectrophotometry

EE – Ethanolic Extract

Declarations

Ethics approval and consent to participate: NIL

Consent for publication: We, the authors have given consent for publication.

Availability of data and materials: Yes, at the request of the authors

Competing interests: No competing interest

Funding: No funding

Authors' contributions: Author 1: Conception, design, and development of the topic, data collection, analysis of data, initial drafting and reviewing the manuscript and final approval of the prepared manuscript. Author 2: Conception, design, and development of research protocol. Author 3: supervision of the experiments, data analysis, and review of the manuscript. Author 4: Student who actually played a supporting role in the experiments and reviewing the manuscript.

Acknowledgements: The authors sincerely acknowledge Akwa Ibom State University for Providing an ambience environment for the research.

5.0 REFERENCES

- Abirami, P. and Rajendran, A. (2011). GC-MS determination of bioactive compounds of *Indigofera aspalathoides*. *J. Nat Prod Plant Resour.*;1(4):126–130.
- Abirami, P. and Rajendran, A. (2012). GC-MS analysis of methanol extracts of *Vernonia cinerea*. *European Journal of Experimental Biology*; 2 (1):9-12.
- Anis, B. N. (2012). Phytochemical composition and antimicrobial activities of essential oil and organic extract from *Pelargonium graveolens* growing in Tunisia. *Lipids in Health and Diseases*; (11):167-171.
- Anwana, E. D., Nyah, G. and Mbong, E. O. (2015). Valuation of Social Relevance of plants amongst the Annangs in South-South Nigeria. *European Journal of Biomedical and Pharmaceuticlas Sciences*; 2:328-337.
- Barreto, F.S.; Sousa, E.O.; Rodrigues, F.F.G.; Costa, J.G.M. and Campos, A. R. (2010). Antibacterial Activity of *Lantana camara* Linn *Lantana montevidensis* Brig Extracts from Cariri-Ceara, Brazil. *J. Young Pharm*; 2: 42–44.
- Begum, S.; Ayub, A.; Qamar Zehra, S.; Shaheen Siddiqui, B.; Iqbal Choudhary, M. Leishmanicidal (2014) Triterpenes from *Lantana camara*. *Chem. Biodivers.*, 11, 709–718.
- Binda, T. K. and Udayan, P. S. (2018). GC-MS analysis of bioactive compounds in methanolic extract of tubers of *Pueraria tuberosa* (Roxb. ex Willd.) DC. - Fabaceae *International Journal of Environment, Agriculture and Biotechnology*; 3(4):1493-1498.
- Clint, C. and Cesar, G. D. (2018). Antimicrobial and cellular metabolic effects of the ethanolic extract of the dallas red variety of *Lantana camara*. *Pharmacophore*; 9(1):10-18.
- Debendranath, M. and Sunita, P. (2020). GC-MS Analysed Phyto-Chemicals and Antibacterial Activity of *WithaniaSomnifera* (L.) Dunal Extract in the Context of Treatment to Liver Cirrhosis. *Biomedical & Pharmacology Journal*; 13(1):71-78.
- Ezekwe, A. S., Rizwan, A. A., Karimah, M. R. and Ewa, O. (2020). Qualitative phytochemical and GC-MS analysis of some commonly consumed vegetables. *GSC Biological and Pharmaceutical Sciences*; 12(03):208-214.
- Ezhilan, B. P. and Neelamegam, R. (2012). GC-MS analysis of phytocomponents in the ethanol extract of *Polygonum chinense* L. *Pharmacogn Res.*; 4(1):11–14.
- Franklin, U., Ohikhen, O. Abosede, W. and Anthony, J. A. (2018). Quantitative Phytochemical Constituents and Antioxidant Activities of the Mistletoe, *Phragmanthera capitata* (Sprengel) Balle Extracted with Different Solvents. *Pharmacognosy Research*; 10(1):16-23.
- Franklin, U., Ohikhen, O., Abosede, W. and Anthony, J. A. (2017). Evaluation of the Antibacterial and Antifungal Properties of *Phragmanthera capitata* (Sprengel) Balle (Loranthaceae), a Mistletoe Growing on Rubber Tree, Using the Dilution Techniques. *The Scientific World Journal*, 1-8.
- George, U. U., Otoh, A. J., Ajayi, O. O. and George, I. E. (2023). Dose-Response Relationship and Histo-morphological Alterations on *Oreochromis niloticus* Juveniles Following Exposure to Ethanolic Extract of *Latana camara*. *Asian Journal of Research in Zoology*; 6(4):71-83.
- George, U. U., Ajayi, O. O., George, I. E. and Vincent, U. E. (2023). Establishing a Dose-response Toxicity for *Clarias gariepinus* Fingerlings Exposed to Ethanolic Extract of *Lantana camara*. *Asian Journal of Fisheries and Aquatic Research*; 24(1):1-10.

16. George, U. U., Otoh, A. J., Ajayi, O. O. and George, I. E. (2023). Studies on Mortality and Histopathological Alteration on the Gills of *Oreochromis niloticus* Juveniles Following Exposure to Ethanolic Extract of *Phragmites capitata* under Laboratory Conditions. *Asian Journal of Fisheries and Aquatic Research*; 24(3):23-34.
17. George, U. U., Mbong, E. O., Abiaobo, N. O. and Akpan, I. I. (2023) In vivo studies on mortality and Histopathological indices of *Phragmites capitata* (Mistletoes) on *Clarias gariepinus* Fingerlings in Aquarium. *Mathews Journal of Cytology and Histology*, 7(2): 1-10.
18. Girish, K. (2017). Antimicrobial Activities of *Lantana Camara* Linn. *Asian J Pharm Clin Res*; 10(3):57-67.
19. Govindaraj, S. and Rajangam, U. (2017). GC-MS analysis of methanolic leaf extracts and stem of *Marsilea minuta* (Linn.) *Journal of Complementary and Alternative Medicine*; 3(1):1-13.
20. Gowri, S. S. (2020). GC-MS analysis of *Majideazanquebarica* Krikex Oliv. (Sapindaceae) seed extract. *Journal of Pharmacological Science & Resources*; 12(5):619-623.
21. Grover, N and Patni, V. (2013). Phytochemical characterization using various solvent extracts and GC-MS analysis of methanolic extract of *Woodfordia fruticosa* (L.) Kurz. Leaves. *Int. J. Phar. & Pharma. Sci.*, 5(4):291-295.
22. Hidayat, R. and Wulandari, P. (2021). Methods of Extraction: Maceration, percolation and Decoction. *Eureka Herba Indonesia*; 2(1):68-74.
23. Kai, L., Xumin, Z., Long, X., Mao, D., Huijuan, C., Jiawen, S., Jiaying, L., Xiaofang, L. and Jia, L. (2020). Lupeol and its derivatives as anticancer and anti-inflammatory agents: Molecular mechanisms and therapeutic efficacy. *Pharmacological Research*; 164(10):53-73.
24. Karimi, E. and Jaafar, H. Z. E. (2011). HPLC and GC-MS determination of bioactive compounds in microwave obtained extracts of three varieties of *Labisia pumila* benth. *Molecules*; 16: 67-71.
25. Kavitha, R. (2021). Phytochemical screening and gc-ms analysis of bioactive compounds present in ethanolic extracts of leaf and fruit of *Trichosanthes dioica* Roxb. *International Journal of Pharmaceutical Sciences and Research*; 12(5):2755-2764.
26. Mariam, I. G., Nouran, M. F., Fulin, W., Maha, M. S., Omar, M. K., Hesham, R. E., Michal, K., Tsong-Long, H., Ahmed, K. O., Mohamed, E., and Shaimaa, F. (2022). Comparative LC-LTQ-MS-MS Analysis of the Leaf Extracts of *Lantana camara* and *Lantana montevidensis* Growing in Egypt with Insights into Their Antioxidant, Anti-Inflammatory, and Cytotoxic Activities. *Plants*; 11:2-21.
27. Naoko, T., Akiko, S., Miki, N., Keisuke, M., Kazutoyo, E., Hajime, H. and Yoshihiro, I. (2007). Antibacterial Activity of Long Chain Fatty Alcohols against *Staphylococcus aureus*. *Molecules*; 12(2): 139-148.
28. Oboho, D. E., Nelson, A. U., Edeke, A., Okwor, J. I., Imakwu, C. A., Akpan, A. U., Oyebadejo, S. and Eyo, J. E. (2022). Phytochemical Screening, GC-MS and Histological Effects of Methanolic Leaf Extract of *Hippocratea Africana* (Willd) on the Midgut of *Sitophilus zeamais* (Motsch). *London Journal of Research in Science*; 22(4):11-22.
29. Pimentel, D., Lach, L., Zuniga, R. and Morrison, D. (2000). Environmental and economic costs of non-indigenous species in the United States. *Biosciences*; 50:53-65.
30. Pinak, P. N. C. and Sudip, C. (2022). GC-MS profiling and bioactivity prediction of compounds from *Momordica charantia* L. extract. *Bioinformation*; 18(10): 1009-1015.
31. Roberts, L. S. and Janovy, J. (2009). Gerald D. Schmidt and Larry S. Roberts' Foundations of Parasitology. McGraw Hill, Boston, 659 Pp.
32. Saliha, R., Syed, T. R., Faizal, A., Absar, A., Shania, A. and Farzana, M. (2014). The Role of Vitamin E in Human Health and some Diseases. *Sultan Qaboos University Medical Journal*; 14(2):157-165.
33. Sathish, R., Vyawahare, B., Natarajan, K. (2011). Antiulcerogenic activity of *Lantana camara* leaves on gastric and duodenal ulcers in experimental rats. *Journal of ethnopharmacology*; 134:195-197.
34. Shettima, A. Y., 1Karumi, Y., 1Sodipo O. A., Usman, H. and Tijjani, M. A. (2013). Gas Chromatography-Mass Spectrometry (GC-MS) Analysis of Bioactive Components of Ethyl Acetate Root Extract of *Guiera senegalensis* J.F. Gmel. *Journal of Applied Pharmaceutical Science*; 3 (3):146-150.
35. Sofowora, A. (2007). Research on Medicinal Plants and Traditional Medicine in Africa. *The Journal of Alternative and Complementary Medicine*; 2(3): 365-372.
36. Tao, X., Siddharth, K. T., Qin, F., Michael, C. L., Marsha, A. W., Melissa, R. J., Melanie, M. M., Scott, R. B., Xing-Cong L., Alice, M. C. and Ameeta, K. A. (2012). A Potent Plant-Derived Antifungal Acetylenic Acid Mediates Its Activity by Interfering with Fatty Acid Homeostasis. *Antimicrobial Agents and Chemotherapy*; 56(6):2894-2907.
37. Tapiero, H., Tew, K. D., Nguyen, B. G. and Mathé, G. (2002). Polyphenols, do they play a role in the prevention of human pathologies? *Biomed. Pharmacother*; 56:200-207.
38. Tin-Lun, Y., Chien-Ming, H., Ling-Jei, M., Tsuimin, T. and Chin-Tin, C. (2022). Oleic Acid-Based Self Micro-Emulsifying Delivery System for Enhancing Antifungal Activities of Clotrimazole. *Pharmaceutics*; 14(3): 1-13.