

Opuntia Cochineal Scale Pest, *Dactylopius opuntiae*: Current Situation, Description, Importance, and Control Methods in Countries of the Mediterranean Basin and North Africa

Mohammed Alhady Abusharya^{ID} and Hosam Ali Ashokri*^{ID}

Zoology Division, Biology Department, Faculty of Science, El-Mergib University, Al-Khums, Libya

Citation: Mohammed Alhady Abusharya and Hosam Ali Ashokri (2025). Opuntia Cochineal Scale Pest, *Dactylopius opuntiae*: Current Situation, Description, Importance, and Control Methods in Countries of the Mediterranean Basin and North Africa. *Acta Biology Forum*. DOI: <https://doi.org/10.51470/ABF.2025.4.2.15>

Corresponding Author: Hosam Ali Ashokri | E-Mail: haalshukri@elmergib.edu.ly

Received 28 April 2025 | Revised 19 May 2025 | Accepted 20 June 2025 | Available Online 25 July 2025

ABSTRACT

This study aims to describe and understand the importance and methods of controlling the cochineal scale insect *Dactylopius opuntiae* (Cockerell, 1896), which has recently spread to most countries in the Mediterranean Basin and North Africa, as well as its original habitats in Mexico and North America. The insect attacks approximately 16 species of cacti, particularly the prickly pear *Opuntia ficus-indica*. Females and nymphs feed on the sap of the plant's stems and branches, causing the stems to dry out and die in severe cases, while males cause little damage. *D. opuntiae* protects itself with a waxy, cotton-like membrane and the secretion of carminic acid (carmine dye). Despite the damage caused by the insect, it is possible to benefit from the carminic acid it secretes, which has antioxidant properties and is used in various medical, food, and commercial applications. Cochineal pest is typically controlled through several methods, including mechanical, biological, and chemical approaches.

Keywords: Cochineal Pest; *Dactylopius opuntiae*; Prickly Pear; Mediterranean Basin; Control Methods.

1. Introduction

The cochineal scale insect *Dactylopius opuntiae* (Cockerell, 1896) is one of the most significant invasive pests, causing substantial economic damage to the Cactaceae family, particularly the prickly pear genus, in various regions of the world [33]. The Cactaceae family is adapted to dry and harsh environments, characterized by high temperatures and low-fertility soils. Prickly pear has a high nutritional value in its fruits, and it is also used as livestock feed and as a fence between farms [36]. The insect is characterized by its thick, waxy covering, which protects it from natural enemies. The carminic acid it secretes helps reduce parasites and predators [25], which has contributed to its widespread distribution in more than 20 countries [16].


The genus *Dactylopius* comprises 11 species, and *Dactylopius opuntiae*, commonly known as the cochineal, is considered one of the most damaging species to cacti [37]. Carminic acid is extracted from the species *Dactylopius coccus* [9], which has numerous medical, food, and commercial applications due to its antioxidant properties [3]. This research article aims to provide a description and definition of this pest, the damage it causes, and how to transform it from a harmful insect to an insect of economic importance.

1.2. General Description & Scientific Classification:

Cochineal (Figure 1) is a small insect belonging to the scale insect family, soft and oval, living in colonies. The colony consists of individuals in different stages. Its males are winged and able to fly.

unlike the females, which are attached to the prickly pear stems

and covered with a white, waxy layer resembling cotton, which acts as a mechanical barrier against external factors such as climate and predators. Females produce a dark purple liquid containing the pigment carmine, which is rich in carminic acid. Females and nymphs feed on the sap in the stems, trunks, and fruits of prickly pear, which, in advanced and severe cases of infection, leads to drying, death, and complete fall of the trunk. As for males, they do not cause any harm to the plant [1]. Based on [2], the female ranges in length from 4 to 6 mm, width from 3 to 4.5 mm, height from 3.8 to 4 mm, and weight from 40 to 47 mg. The ventral and dorsal appearance of the female is cylindrical, the antennae are composed of seven segments, the breathing holes are large, and the legs are short and strong [4].



Cochineal pest was first described by Cockerell in 1896. It belongs to the order Hemiptera within the family Dactylopiidae and is known by the scientific name *Dactylopius opuntiae* [10].

Kingdom: Animalia
Phylum: Arthropoda
Class: Insecta
Order: Hemiptera
Suborder: Sternorrhyncha
Family: Dactylopiidae
Genus: *Dactylopius*
Species: *opuntiae*

Figure 1: Female (Left) & Male (Right) Cochineal Drawing.

Copyright: This is an open-access article distributed under the terms of the Creative Commons Attribution License (CC BY 4.0), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

1.3. Life Cycle:

As illustrated in Figure 2, the female goes through three stages of development: the egg, the nymphal stage which consists of two stages (the first nymphal instar and the second nymphal instar), and the adult insect stage, while the male goes through the following stages: the egg, the nymphal stage, the pre-pupa, the pupa, and the adult insect which carries a pair of wings [2]. The eggs are bright red, oval in shape, and have a smooth surface. The average length of the egg is 0.7 mm, and its width is 0.3 mm [24].




Figure 2: Lifecycle of *Dactylopius opuntiae* [19]

The female lays eggs individually or in groups, linked in a chain [30]. The eggs hatch approximately 18-27 minutes after laying. The female lays approximately 226 eggs during her lifetime. The average duration of one generation ranges from 56 to 58 days from egg laying until the adult insect emerges. The life cycle of females ranges from 40-180 days, while that of males does not exceed 35-52 days, depending on climatic conditions [27]; [28]. *D. opuntiae* usually reproduces sexually, but may tend to reproduce parthenogenetically to produce fewer offspring under harsh conditions, especially during high temperatures [14]; [32]. The insect produces between 4 and 5 generations per year, especially in warm regions [27].

1.4. Means of insect spread:

According to [17], the high fertility and the waxy cover that protects the insect are factors that have helped it spread in several environments. This insect is also easily spread by wind, rain, and water, in addition to its spread by birds, humans, and livestock.

1.5. Damage and symptoms:

The cochineal *D. opuntiae* is a harmful insect that infects about 16 species of cactus worldwide, causing significant losses, especially to the prickly pear genus and its various species [25]; [33]. As shown in Figure 3, the plant first appears as yellow-green areas and necrosis on the branches, which gradually expand, then dry up within about a year [25]; [11], while the woody stems can remain alive for another 6 months after the branches die and dry up [18]. Then, the large colonies of *D. opuntiae* increase the growth of endophytic fungi within the host and host plant as a result of stem and trunk rot [15].



Figure 3: Prickly Pear Infected with Cochineal

1.6. World Geographical Distribution:

The native habitat of *Dactylopius opuntiae* is Mexico, and the current distribution of this insect includes many countries around the world. Based on [16], the most important countries in which the insect has been recorded are: Australia, Brazil, Hawaii, India, Jamaica, Madagascar, Mexico, South Africa, Sri Lanka, the United States, Zimbabwe, Pakistan, France, Spain, India, and Cyprus. The presence of *D. Opuntiae* has also been recorded in Arab countries, including Morocco, Lebanon, Palestine, Saudi Arabia, Jordan, Syria, as well as Yemen. A leading study conducted by [20] reported the presence of the cochineal insect in the state of Algeria in the city of Misserghin (Oran) in 2024. In Libya, the insect first appeared in the western coastal areas such as Zawiya and Abu Surra in May 2024, then spread to other areas such as JudDaim, Al-Maya, Al-Tuwaibiya, Al-Maamura, Qarquza, and Al-Zahra. It is believed that it spread from neighboring countries via wind, agricultural machinery, sheep, or fodder. The insect feeds exclusively on the sap of the prickly pear plant, causing the pads to yellow and fall off, turn brown, and eventually die if left uncontrolled.




Figure 4: World Distribution of *Dactylopius opuntiae* [38]

1.7. Industrial and Economic Importance:

The cochineal insect *D. opuntiae* has long been used in biological control to control the spread of several cactus species, such as *Opuntia stricta* and *O. ficus-indica* [27]. Carminic acid extracted from the insect has antioxidant properties and has been used in beef storage and preservation [3]. Carminic acid also helps inhibit tumors and reduce the progression of fatty liver disease [21].

The red dye carmine is widely used in cosmetics, pharmaceuticals, food, and the clothing industry [11]. Due to the insect's reproductive capacity and the damage it causes to cactus stems, this pest is considered one of the pests that cause significant economic and agricultural losses in several countries around the world [35].

1.8. Cochineal Control Techniques:

Given the widespread damage caused by the insect, methods are required to limit its spread. These methods consist of several techniques based on mechanical, physical, biological, chemical, and other methods, which can be combined in various ways to achieve the best results in controlling the pest.

1.8.1 Mechanical and Physical Methods: Mechanical methods are highly effective, especially in areas where the first outbreak of *D. opuntiae* is detected. This is achieved through regular and continuous field visits. This method mainly consists of harvesting or mechanically cleaning infected plant parts and destroying them by burning or burying them as quickly as possible. It is also possible to preserve and store them for use as livestock feed [34].

1.8.2 Biological Control Methods: Many natural enemies of *D. opuntiae* have been recorded, some of which are capable of controlling the insect. These predators include insects and spiders. The most important predators of *D. opuntiae* are the beetles belonging to the order Coleoptera: *Hyperaspis trifurcata*, *Chilocorus cacti*, *Coccidophilus citricola*, *Zagreus bimaculosus*, *Exochomus flavipes*, *Exochomus flaviventris*, *Cryptolaemus montrouzieri*, and *Cybocephalus* sp[22]; [31]. The beetle *Hyperaspis trifurcata* is considered the most efficient and effective due to its abundance [8]. This beetle is used as a biological control agent in many ecosystems and agricultural systems due to its strong predatory activity, as the female consumes more than 5,400 nymphs during her lifetime, making it one of the most important predators of the cochineal insect *D. opuntiae*[35]. Several insects also feed on the nymphal stage of *D. opuntiae*, including the aphid *Leucopis bellula* and the brown barber worm *Sympherobius barberi*[29], while the snout moth *Laetilia coccidivora* feeds on the adult stage of *D. opuntiae*[26]; [13].

1.8.3 Chemical Control Methods: In the early stages of chemical control, *D. opuntiae* colonies are sprayed and treated with insecticides and some detergents [12]; [23]. When the prevalence of *D. opuntiae* increases significantly and it is difficult to eliminate it using biological and mechanical methods, chemical compounds are used. These compounds are mainly used in Mexico, which relies on malathion, methyl parathion, and chlorpyrifos [8], and triclofen[5]. Cypermethrin, carbaryl, and chlorpyrifos have been used in South Africa [25]. The use of mineral oil has been successful in eliminating *D. opuntiae* by 93–97% [7] and 100% [6].

2. Conclusion:

The insect of *D. opuntiae* causes severe damage to cactus species worldwide. It feeds on plant sap, causing the plant to desiccate and die. It is widespread in several countries. The female produces carminic acid, which has antioxidant properties, making it an important ingredient in many industries and economically important. The insect has also been used as a pest control for cactus plants in some countries.

Mechanical, biological, and chemical control methods are among the most important.

3. Recommendations:

The Authors of this work would highly recommend an immediate action plan that includes:

- Launching control campaigns in affected areas. These campaigns aim to contain the spread of the pest and prevent its transmission to new areas.
- Continuous monitoring and evaluation of the results of the control campaigns.
- Launching national awareness campaigns for farmers about the risks of this pest and the importance of early detection and immediate reporting.

References

1. Al-Hindi, A.N., Al-Hawshabi, O.S., Bawazer, G.M., Busais, S.M., Baeshen, O.A., Al Khasha, M.S., Bin Othman, A., Awlaki, W., Ismail Hassan, I., Metwally, F.(2022). A new record of the cochineal insect (*Dactylopius opuntiae*) on the prickly pear (*Opuntia dellenii*) in Yemen - tur al-Baha - lahj. *Electronic Journal of University of Aden for Basic and Applied Sciences*, 3: (2): 66-74.
2. Alhousa, W., Basheer, A.N., El-Naser, Z.(2024). Study of the Biology of the Cochineal Scale Insect, *Dactylopius opuntiae*. *Arab Journal of Plant Protection*, 42(2): 143-148.
3. Aragon-Martinez, O.H., Martinez-Morales, F., González-Chávez, M.M., Méndez-Gallegos, S.d.J., González-Chávez, R., Posadas-Hurtado, J.C., Isiordia-Espinoza, M.A.(2023). *Dactylopius opuntiae* [Cockerell] Could Be a Source of Antioxidants for the Preservation of Beef Patties. *Insects*, 1-12.
4. Bader, A., & Abu Alloush, A. (2019). First Record of the Cochineal Scale Insect, *Dactylopius opuntiae* (Cockerell) (Hemiptera: Dactylopiidae), in Jordan, *Jordan Journal of Biological Sciences*, 12: 155-159.
5. Badii, M.H., & Flores, A.E. (2001). Prickly pear cacti pests and their control in Mexico. *Florida Entomologist*, 84: 503-505.
6. Borges, L.R., Santos, D.C., Gomes, E.W.F., Cavalcanti, V.A.L.B., Silva, I.M.M., Falcão, H.M. da Silva, D.M.P. (2013). Use of biodegradable products for the control of *Dactylopius opuntiae* (Hemiptera: Dactylopiidae) in cactus pear. *Acta Horticulturae*. 995: 379-386
7. Brito, C.H., Lopes, E.B., Albuquerque, I.C., Luna Batista, J. (2008). Avaliação de produtos alternativos e pesticidas no controle da cochonilha-do-carmimna Paraíba. *Revista de Biologia e Ciências da Terra* 8: 1-5.
8. Cruz-Rodríguez. J.A., González-Machorro. E., Villegas González A.A., Rodríguez Ramírez, M.L., Mejía Lara. F. (2016) Autonomous biological control of *Dactylopius opuntiae* (Hemiptera: Dactylopiidae) in a prickly pear plantation with ecological management. *Environmental Entomology* 45: 642-648.
9. de Carvalho, P.R., Bolognesi, V.J., Longhi, D.A., Rocha Garcia, C.E. (2021). Assessment of the lipid oxidation, color, and sensorial acceptance of fresh sausage formulated with potassium bixinate as a substitute for sodium nitrite and carmine. *J. Food Sci. Technol.* 58, 4524-4529.
10. De Lotto, G. (1974). On the status and identity of the cochineal insects (Homoptera: Coccoidea: Dactylopiidae). *Journal of the Entomological Society of Southern Africa*, 37: 167-193.
11. El Aalaoui, M., Mokri, F., Dababat, A.A., Lahlali, R., Sbaghi, M. (2022). Moroccan entomopathogenic nematodes as potential biocontrol agents against *Dactylopius opuntiae* (Hemiptera: Dactylopiidae). *Sci. Rep.* 12, 7590.
12. Fand, B., Suroshe, S., Gautam, R. (2013). Fortuitous biological control of insect pests and weeds: A critical review. *The Bioscan* 8 (1), 1-10.

13. Flores, A., Olvera, H., Cruz-Rodríguez, S., Barranco, J. (2013). Predation potential of *Chilocorus cacti* (Coleoptera: Coccinellidae) to the prickly pear cacti pest *Dactylopius opuntiae* (Hemiptera: Dactylopiidae). *Neotropical Entomology* 42: 407-411.
14. Flores-Hernández, A., Murillo-Amador, B., Rueda-Puente, E.O., Salazar-Torres, J.C., García-Hernández, J.L., Troyo-Díez, E. (2006). Reproducción de cochinitas silvestre *Dactylopius coccus* (Homóptera: Dactylopiidae). *Revista Mexicana de Biodiversidad*, 77: 97-102.
15. Freire, K.T., Araújo, G.R., Bezerra, J.D.P., Barbosa, R.N., Silva, D.C., Svedese, V., Paiva, L., Souza-Motta, C. (2015). Fungos endofíticos de *Opuntia ficus-indica* (L.) Mill. (Cactaceae) sadio e infestado por *Dactylopius opuntiae* (Cockerell, 1896) (Hemiptera: Dactylopiidae). *Gaia Scientia*, 9: 104-110.
16. García Morales, M., Denno, B.D., Miller, D.R., Miller, G.L., Bendov, Y., Hardy, N.B. (2016). ScaleNet: A literature-based model of scale insect biology and systematics. Database. *The Journal of Biological Databases and Curation*. 2016.
17. Khan, H.A.A., Sayyed, A.H., Akram, W., Raza, S., Ali, M. (2012). The predatory potential of *Chrysoperla carnea* and *Cryptolaemus montrouzieri* larvae on different stages of the mealybug, *Phenacoccus solenopsis*: A threat to cotton in South Asia. *Journal of Insect Science*, 12(1):147. Available online: <http://www.insectscience.org/12.147>
18. Klein, H. (2002). Biological control of invasive cactus species (Family Cactaceae). 2.2. Cochineal insects (*Dactylopius* spp.). ARC-Plant Protection Research Institute, PPRI leaflet series, *Weeds Biocontrol*, 2:1-3.
19. Kondo, T., & Watson, G. W. (2022). *Encyclopedia of scale insect pests*: CABI.
20. Leblalta, A., Meradsi, F., Katbeh Bader, A., Kenou, G. B., Warot, S., Ris, N. (2025). First detection of the opuntia cochineal scale *Dactylopius opuntiae* (Cockerell, 1896) (Hemiptera: Dactylopiidae), in Misserghin (Oran), the North-West of Algeria. *Acta Phytopathologica et Entomologica Hungarica* (published online ahead of print 2025). 10.1556/038.2025.00235.
21. Li, L., Fang, B., Zhang, Y., Yan, L., He, Y., Hu, L., Xu, Q., Li, Q., Dai, X., Kuang, Q., Xu, M., Tanm, J., Ge, C. (2022). Carminic acid mitigates fructose-triggered hepatic steatosis by inhibition of oxidative stress and inflammatory reaction. *Biomed. Pharmacother*, 145.
22. Lima, M.S., Da Silva, D.M.P., Falcão, H.M., Ferreira, W.M., Silva, L.D., Paranhos, B.A.J. (2011). Predadores associados a *Dactylopius opuntiae* (Hemiptera: Dactylopiidae) empalme forrageira no Estado de Pernambuco, Brasil. *Revista Chilena de Entomología*, 36: 51-54.
23. López-Rodríguez, P.E., Aquino-Pérez, G., Morales-Flores, F.J., Mena-Covarrubias, J., Rodríguez-Leyva, E., de Jesús Méndez-Gallegos, S. (2021). Productos no convencionales como alternativa de control de *Dactylopius opuntiae* Cockerell (Hemiptera: Dactylopiidae). *Revista Fitotecnia Mexicana*. 44, 417.
24. Mathenge, C.W., Holford, P., Hoffmann, J.H., Spooner-Hart, R., Beattie, G.A.C., Zimmermann, H.G. (2009). The biology of *Dactylopius tomentosus* (Hemiptera: Dactylopiidae). *Bulletin of Entomological Research*, 99(6):551-559.
25. Mazzeo, G., Nucifora, S., Russo, A., Suma, P. (2019). *Dactylopius opuntiae*, a new prickly pear cactus pest in the Mediterranean: An overview. *Entomologia Experimentalis et Applicata*, 167(1):59-72.
26. Mengual, X., & Thompson, C. (2011). Carmine cochineal killers: the flower fly genus *Eosalpingogaster* Hull (Diptera: Syrphidae) revised. *Systematic Entomology* 36: 713-731.
27. Moussa, Z., Yammouni, D., Azar, D. (2017). *Dactylopius opuntiae* (Cockerell, 1896), a new invasive pest of the cactus plants *Opuntia ficus-indica* in the South of Lebanon (Hemiptera, Coccoidea, Dactylopiidae). *Bull Soc Entomol France*, 122(2): 173-178.
28. Ochoa, M.J., Lobos, E., Portillo, L., Vigueras, A.L. (2015). Importance of biotic factors and impact on cactus pear production systems. *Acta Horticulture*, 1067: 327-333.
29. Pacheco Rueda, I., Lomeli Flores, J.R., Rodríguez Leyva, E., Ramírez Delgado, M. (2011). Ciclo de vida y parámetros poblacionales de *Symphorobius barberi* Banks (Neuroptera: Hemerobiidae) criado con *Dactylopius opuntiae* Cockerell (Hemiptera: Dactylopiidae). *Acta Zoológica Mexicana*, 27: 325-340.
30. Palafox-Luna, A., Rodríguez-Leyva, E., Lomeli-Flores, R., Vigueras-Guzmán, L., Vanegas-Rico, M. (2018). Life cycle and fecundity of *Dactylopius opuntiae* (Hemiptera: Dactylopiidae) in *Opuntia ficus-indica* (Caryophyllales: Cactaceae). *Agrociencia*, 52(1):103-114
31. Ramírez, S.O., Cruz-Rodríguez, J.A., Gamero, A.M. (2011). Enemigos naturales de la cochinita silvestre, *Dactylopius* spp. (Hemiptera: Dactylopiidae), en un huerto orgánico de nopal tunero, en México. Workshop, Agroecología e Desenvolvimento Sustentável (ed. by A Bento, JA Pereira & MA Rodrigues), pp. 5-13. *Escola Superior Agrária de Bragança*, Bragança, Portugal.
32. Romero López, B.E., Flores Hernández, A., Santamaría César, E., Salazar Torres, J.C., Ramírez Delgado, M., Pedroza Sandoval, A. (2006). Identificación, biología y adaptación de la cochinita silvestre *Dactylopius opuntiae* (Homóptera: Dactylopiidae) a las condiciones ambientales de bermejillo, durango.. *Revista Chapingo Serie Zonas Áridas*, V (1), 41-48.
33. Rule, N.F., & Hoffmann, J. (2018). "The performance of *Dactylopius opuntiae* as a biological control agent on two invasive *Opuntia* cactus species in South Africa". *Biological Control*, 119: 7-11.
34. Torres, J.B., & Giorgi, J.A. (2018). Management of the false carmine cochineal *Dactylopius opuntiae* (Cockerell): perspective from Pernambuco state, Brazil. *Phytoparasitica*, 46: 331-340.
35. Vanegas-Rico, J.M., Rodríguez-Leyva, E., Lomeli-Flores, J.R., Gonzalez-Hernandez, H., Perez-Panduro, A., Mora-Aguilera, G. (2016). Biology and life history of *Hyperaspis trifurcata* feeding on *Dactylopius opuntiae*. *BioControl*, 61: 691-701.
36. Vastolo, A., Calabrò, S., Cutrignelli, M.I., Raso, G., Todaro, M. (2020). Silage of Prickly Pears (*Opuntia*) juice By-products. *Animals*. 10.1716.
37. Vera-Ponce León, A., Dominguez-Mirazo, M., Bustamante-Brito, R., Higareda Alvear, V., Rosenbluth, M., Martínez-Romero, E. (2021). Functional genomics of a *Spiroplasma* associated with the carmine cochineals *Dactylopius coccus* and *Dactylopius opuntiae*. *BMC Genom*, 22, 240.
38. Zahra, S., Sammut, M., Debono, J. (2018). Phytobezoar large bowel obstruction-The prickly pear (A single centre experience and case series report). *International Journal of Surgery Open*, 13, 10-14.